

## UNIFORM GLOBAL ATTRACTORS FOR NON-AUTONOMOUS DISSIPATIVE DYNAMICAL SYSTEMS

MICHAEL ZGUROVSKY

National Technical University of Ukraine “Igor Sikorsky Kyiv Polytechnic Institute”  
Peremogy ave., 37, build, 1, 03056, Kyiv, Ukraine

MARK GLUZMAN

Department of Applied Physics and Applied Mathematics, Columbia University  
New York, NY 10027, USA

NATALIIA GORBAN, PAVLO KASYANOV \*  
LILIIA PALIICHUK AND OLHA KHOMENKO

Institute for Applied System Analysis  
National Technical University of Ukraine “Igor Sikorsky Kyiv Polytechnic Institute”  
Peremogy ave., 37, build, 35, 03056, Kyiv, Ukraine

**ABSTRACT.** In this paper we consider sufficient conditions for the existence of uniform compact global attractor for non-autonomous dynamical systems in special classes of infinite-dimensional phase spaces. The obtained generalizations allow us to avoid the restrictive compactness assumptions on the space of shifts of non-autonomous terms in particular evolution problems. The results are applied to several evolution inclusions.

**1. Introduction.** The standard scheme of investigation of uniform the long-time behavior for all solutions of non-autonomous problems covers non-autonomous problems of the form

$$\partial_t u(t) = A_{\sigma(t)}(u(t)), \quad (1)$$

where  $\sigma(s)$ ,  $s \geq 0$ , is a functional parameter called the time symbol of equation (1) ( $t$  is replaced by  $s$ ). In applications to mathematical physics equations, a function  $\sigma(s)$  consists of all time-dependent terms of the equation under consideration: external forces, parameters of mediums, interaction functions, control functions, etc; Chepyzhov and Vishik [4, 5, 8]; Sell [36]; Zgurovsky et al. [46] and references therein; see also Hale [16]; Ladyzhenskaya [30]; Mel’nik and Valero [32]; Iovane, Kapustyan and Valero [17]. In the mentioned above papers and books it is assumed that the symbol  $\sigma$  of equation (1) belongs to a Hausdorff topological space  $\Xi_+$  of functions defined on  $\mathbb{R}_+$  with values in some complete metric space. Usually, in applications, the topology in the space  $\Xi_+$  is a local convergence topology on any segment  $[t_1, t_2] \subset \mathbb{R}_+$ . Further, they consider the family of equations (1) with various symbols  $\sigma(s)$  belonging to a set  $\Sigma \subseteq \Xi_+$ . The set  $\Sigma$  is called the symbol space

---

2010 *Mathematics Subject Classification.* Primary: 37L05, 35B41; Secondary: 35B40.

*Key words and phrases.* Global attractor, multivalued semi-flow, non-autonomous dynamical system, evolution inclusion, long-time behavior.

The research was partially supported by the National Academy of Sciences of Ukraine under grant 2284/15 and by Grant of the President of Ukraine GP/F61/017.

\* Corresponding author: Pavlo Kasyanov.

of the family of equations (1). It is assumed that the set  $\Sigma$ , together with any symbol  $\sigma(s) \in \Sigma$ , contains all positive translations of  $\sigma(s)$ :  $\sigma(t+s) = T(t)\sigma(s) \in \Sigma$  for any  $t, s \geq 0$ . The symbol space  $\Sigma$  is invariant with respect to the translation semigroup  $\{T(t)\}_{t \geq 0}$ :  $T(t)\Sigma \subseteq \Sigma$  for any  $t \geq 0$ . To prove the existence of uniform trajectory attractors they suppose that the symbol space  $\Sigma$  with the topology induced from  $\Xi_+$  is a compact metric space. Mostly in applications, as a symbol space  $\Sigma$  it is naturally to consider the hull of translation-compact function  $\sigma_0(s)$  in an appropriate Hausdorff topological space  $\Xi_+$ . The direct realization of this approach to differential-operator inclusions, PDEs with Caratheodory's nonlinearities, optimization problems, etc, is problematic without any additional assumptions for parameters of Problem (1) and requires the translation-compactness of the symbol  $\sigma(s)$  in some compact Hausdorff topological space of measurable multivalued mappings acts from  $\mathbb{R}_+$  to some metric space of operators from  $(V \rightarrow 2^{V^*})$ , where  $V$  is a Banach space and  $V^*$  is its dual space, satisfying (possibly) only growth and sign assumptions. To avoid this technical difficulties we present an alternative approach for the existence and construction of the uniform global attractor for classes of non-autonomous dynamical systems in special classes of infinite-dimensional phase spaces.

**2. Main constructions and results.** Let  $p \geq 2$  and  $q > 1$  be such that  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $(V; H; V^*)$  to be evolution triple such that  $V \subset H$  with compact embedding. For each  $t_1, t_2 \in \mathbb{R}$ ,  $0 \leq t_1 < t_2 < +\infty$ , consider the space

$$W_{t_1, t_2} := \{y(\cdot) \in L_p(t_1, t_2; V) : y'(\cdot) \in L_q(t_1, t_2; V^*)\},$$

where  $y'(\cdot)$  is a derivative of an element  $y(\cdot) \in L_p(t_1, t_2; V)$  in the sense of distributions  $\mathcal{D}^*([t_1, t_2]; V^*)$ . The space  $W_{t_1, t_2}$  endowed with the norm

$$\|y\|_{W_{t_1, t_2}} := \|y\|_{L_p(t_1, t_2; V)} + \|y'\|_{L_q(t_1, t_2; V^*)}, \quad y \in W_{t_1, t_2},$$

is a reflexive Banach space. Note that  $W_{t_1, t_2} \subset C([t_1, t_2]; H)$  with continuous and dense embedding; Gajewsky et al [11, Chapter IV]. For each  $\tau \geq 0$ , consider the Fréchet space

$$W^{\text{loc}}([\tau, +\infty)) := \{y : [\tau, +\infty) \rightarrow H : \Pi_{t_1, t_2} y \in W_{t_1, t_2} \text{ for each } [t_1, t_2] \subset [\tau, +\infty)\},$$

where  $\Pi_{t_1, t_2}$  is the restriction operator to the finite time interval  $[t_1, t_2]$ . We recall that the sequence  $\{f_n\}_{n \geq 1}$  converges in  $W^{\text{loc}}([\tau, +\infty))$  (in  $C^{\text{loc}}([\tau, +\infty); H)$  respectively) to  $f \in W^{\text{loc}}([\tau, +\infty))$  (to  $f \in C^{\text{loc}}([\tau, +\infty); H)$  respectively) as  $n \rightarrow +\infty$  if and only if the sequence  $\{\Pi_{t_1, t_2} f_n\}_{n \geq 1}$  converges in  $W_{t_1, t_2}$  (in  $C([t_1, t_2]; H)$  respectively) to  $\Pi_{t_1, t_2} f$  as  $n \rightarrow +\infty$  for each finite time interval  $[t_1, t_2] \subset [\tau, +\infty)$ . Further we denote that

$$T(h)y(\cdot) = \Pi_{0, +\infty} y(\cdot + h), \quad y \in W^{\text{loc}}(\mathbb{R}_+), \quad h \geq 0,$$

where  $\mathbb{R}_+ = [0, +\infty)$  and  $\Pi_{0, +\infty}$  is the restriction operator to the time interval  $[0, +\infty)$ .

Throughout the paper we consider the family of solution sets  $\{\mathcal{K}_\tau^+\}_{\tau \geq 0}$  such that  $\mathcal{K}_\tau^+ \subset W^{\text{loc}}([\tau, +\infty))$  for each  $\tau \geq 0$  and  $\mathcal{K}_{\tau_0}^+ \neq \emptyset$  for some  $\tau_0 \geq 0$ . In the most of applications as  $\mathcal{K}_\tau^+$  can be considered the family of globally defined on  $[\tau, +\infty)$  weak solutions for particular non-autonomous evolution problem (see Section 4).

To state the main assumptions on the family of solution sets  $\{\mathcal{K}_\tau^+\}_{\tau \geq 0}$  it is necessary to formulate two auxiliary definitions.

A function  $\varphi \in L^\gamma_{\text{loc}}(\mathbb{R}_+)$ ,  $\gamma > 1$ , is called *translation bounded function* in  $L^\gamma_{\text{loc}}(\mathbb{R}_+)$  if

$$\sup_{t \geq 0} \int_t^{t+1} |\varphi(s)|^\gamma ds < +\infty;$$

Chepyzhov and Vishik [7, p. 105]. A function  $\varphi \in L^1_{\text{loc}}(\mathbb{R}_+)$  is called a *translation uniform integrable (t.u.i.) function* in  $L^1_{\text{loc}}(\mathbb{R}_+)$  if

$$\lim_{K \rightarrow +\infty} \sup_{t \geq 0} \int_t^{t+1} |\varphi(s)| \mathbf{I}\{|\varphi(s)| \geq K\} ds = 0;$$

Gorban et al. [13]. Note that Dunford-Pettis compactness criterion provides that  $\varphi \in L^1_{\text{loc}}(\mathbb{R}_+)$  is a t.u.i. function in  $L^1_{\text{loc}}(\mathbb{R}_+)$  if and only if for every sequence of elements  $\{\tau_n\}_{n \geq 1} \subset \mathbb{R}_+$ , the sequence  $\{\varphi(\cdot + \tau_n)\}_{n \geq 1}$  contains a subsequence converging weakly in  $L^1_{\text{loc}}(\mathbb{R}_+)$ . Note that for each  $\gamma > 1$ , every translation bounded in  $L^\gamma_{\text{loc}}(\mathbb{R}_+)$  function is t.u.i. in  $L^1_{\text{loc}}(\mathbb{R}_+)$ ; Gorban et al. [13].

**Main assumptions.** Let the following two assumptions hold:

(A1) there exist a t.u.i. in  $L^1_{\text{loc}}(\mathbb{R}_+)$  function  $c_1 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  and a constant  $\alpha_1 > 0$  such that for each  $\tau \geq 0$ ,  $y \in \mathcal{K}^+_\tau$ , and  $t_2 \geq t_1 \geq \tau$ , the following inequality holds:

$$\|y(t_2)\|_H^2 - \|y(t_1)\|_H^2 + \alpha_1 \int_{t_1}^{t_2} \|y(t)\|_V^p dt \leq \int_{t_1}^{t_2} c_1(t) dt; \tag{2}$$

(A2) there exist a t.u.i. in  $L^1_{\text{loc}}(\mathbb{R}_+)$  function  $c_2 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  and a constant  $\alpha_2 > 0$  such that for each  $\tau \geq 0$ ,  $y \in \mathcal{K}^+_\tau$ , and  $t_2 \geq t_1 \geq \tau$ , the following inequality holds:

$$\int_{t_1}^{t_2} \|y'(t)\|_{V^*}^q dt \leq \alpha_2 \int_{t_1}^{t_2} \|y(t)\|_V^p dt + \int_{t_1}^{t_2} c_2(t) dt. \tag{3}$$

To characterize the uniform long-time behavior of solutions for non-autonomous dissipative dynamical system consider the *united trajectory space*  $\mathcal{K}^+_{\cup}$  for the family of solutions  $\{\mathcal{K}^+_\tau\}_{\tau \geq 0}$  shifted to zero:

$$\mathcal{K}^+_{\cup} := \bigcup_{\tau \geq 0} \{T(h)y(\cdot + \tau) : y(\cdot) \in \mathcal{K}^+_\tau, h \geq 0\} \subset W^{\text{loc}}(\mathbb{R}_+), \tag{4}$$

and the *extended united trajectory space* for the family  $\{\mathcal{K}^+_\tau\}_{\tau \geq 0}$ :

$$\mathcal{K}^+ := \text{cl}_{C^{\text{loc}}(\mathbb{R}_+; H)} [\mathcal{K}^+_{\cup}], \tag{5}$$

where  $\text{cl}_{C^{\text{loc}}(\mathbb{R}_+; H)}[\cdot]$  is the closure in  $C^{\text{loc}}(\mathbb{R}_+; H)$ . Since  $T(h)\mathcal{K}^+_{\cup} \subseteq \mathcal{K}^+_{\cup}$  for each  $h \geq 0$ , then

$$T(h)\mathcal{K}^+ \subseteq \mathcal{K}^+ \text{ for each } h \geq 0, \tag{6}$$

due to

$$\rho_{C^{\text{loc}}(\mathbb{R}_+; H)}(T(h)u, T(h)v) \leq \rho_{C^{\text{loc}}(\mathbb{R}_+; H)}(u, v) \text{ for each } u, v \in C^{\text{loc}}(\mathbb{R}_+; H),$$

where  $\rho_{C^{\text{loc}}(\mathbb{R}_+; H)}$  is the standard metric on Fréchet space  $C^{\text{loc}}(\mathbb{R}_+; H)$ . Therefore the set

$$\mathbb{X} := \{y(0) : y \in \mathcal{K}^+\} \tag{7}$$

is closed in  $H$  (it follows from Theorem 3.1). We endow this set  $\mathbb{X}$  with metric

$$\rho_{\mathbb{X}}(x_1, x_2) = \|x_1 - x_2\|_H, \quad x_1, x_2 \in \mathbb{X}.$$

Then we obtain that  $(\mathbb{X}, \rho)$  is a Polish space (complete separable metric space).

Let us define the multivalued semiflow (*m-semiflow*)  $G : \mathbb{R}_+ \times \mathbb{X} \rightarrow 2^{\mathbb{X}}$ :

$$G(t, y_0) := \{y(t) : y(\cdot) \in \mathcal{K}^+ \text{ and } y(0) = y_0\}, \quad t \geq 0, y_0 \in \mathbb{X}. \tag{8}$$

According to (6), (7), and (8) for each  $t \geq 0$  and  $y_0 \in \mathbb{X}$  the set  $G(t, y_0)$  is nonempty. Moreover, the following two conditions hold:

- (i)  $G(0, \cdot) = I$  is the identity map;
- (ii)  $G(t_1 + t_2, y_0) \subseteq G(t_1, G(t_2, y_0))$ ,  $\forall t_1, t_2 \in \mathbb{R}_+, \forall y_0 \in \mathbb{X}$ ,

where  $G(t, D) = \bigcup_{y \in D} G(t, y)$ ,  $D \subseteq \mathbb{X}$ .

We denote by  $\text{dist}_{\mathbb{X}}(C, D) = \sup_{c \in C} \inf_{d \in D} \rho(c, d)$  the *Hausdorff semidistance* between nonempty subsets  $C$  and  $D$  of the Polish space  $\mathbb{X}$ . Recall that the set  $\mathfrak{R} \subset \mathbb{X}$  is a *global attractor* of the m-semiflow  $G$  if it satisfies the following conditions:

- (i)  $\mathfrak{R}$  attracts each bounded subset  $B \subset \mathbb{X}$ , i.e.

$$\text{dist}_{\mathbb{X}}(G(t, B), \mathfrak{R}) \rightarrow 0, \quad t \rightarrow +\infty; \tag{9}$$

- (ii)  $\mathfrak{R}$  is negatively semi-invariant set, i.e.  $\mathfrak{R} \subseteq G(t, \mathfrak{R})$  for each  $t \geq 0$ ;
- (iii)  $\mathfrak{R}$  is the minimal set among all nonempty closed subsets  $C \subseteq \mathbb{X}$  that satisfy (9).

In this paper we examine the uniform long-time behavior of solution sets  $\{\mathcal{K}_\tau^+\}_{\tau \geq 0}$  in the strong topology of the natural phase space  $H$  (as time  $t \rightarrow +\infty$ ) in the sense of the existence of a compact global attractor for m-semiflow  $G$  generated by the family of solution sets  $\{\mathcal{K}_\tau^+\}_{\tau \geq 0}$  and their shifts. The following theorem is the main result of the paper.

**Theorem 2.1.** *Let assumptions (A1)–(A2) hold. Then the m-semiflow  $G$ , defined in (8), has a compact global attractor  $\mathfrak{R}$  in the phase space  $\mathbb{X}$ .*

**3. Proof of Theorem 2.1.** Before the proof of Theorem 2.1 we provide the following statement characterizing the compactness properties of the family  $\mathcal{K}^+$  in the topology induced from  $C^{loc}(\mathbb{R}_+; H)$ .

**Theorem 3.1.** *Let assumptions (A1)–(A2) hold. Then the following two statements hold:*

- (a) for each  $y \in \mathcal{K}^+$ , the following estimate holds

$$\|y(t)\|_H^2 \leq \|y(0)\|_H^2 e^{-c_3 t} + c_4, \quad t \geq 0, \tag{10}$$

where the positive constants  $c_3$  and  $c_4$  do not depend on  $y \in \mathcal{K}^+$  and  $t \geq 0$ ;

- (b) for any bounded in  $L_\infty(\mathbb{R}_+; H)$  sequence  $\{y_n\}_{n \geq 1} \subset \mathcal{K}^+$ , there exist an increasing sequence  $\{n_k\}_{k \geq 1} \subseteq \mathbb{N}$  and an element  $y \in \mathcal{K}^+$  such that

$$\|\Pi_{\tau, T} y_{n_k} - \Pi_{\tau, T} y\|_{C([\tau, T]; H)} \rightarrow 0, \quad k \rightarrow +\infty, \tag{11}$$

for each finite time interval  $[\tau, T] \subset (0, +\infty)$ . If, additionally, there exists  $y_0 \in H$  such that  $y_{n_k}(0) \rightarrow y_0$  in  $H$ , then  $y(0) = y_0$ .

*Proof of Theorem 3.1.* Let us prove statement (a). If statement (a) holds for each  $y \in \mathcal{K}_\cup^+$ , then inequality (10) holds for each  $y \in \mathcal{K}^+$ , due to (5). The rest of the proof of statement (a) establishes inequality (10) for each  $y \in \mathcal{K}_\cup^+$ .

For an arbitrary  $y \in \mathcal{K}_\cup^+$ , there exist  $\tau, h \geq 0$  and  $z(\cdot) \in \mathcal{K}_\tau^+$  such that  $y(\cdot) = T(\tau + h)z(\cdot)$ . Assumption (A1) implies the following inequality:

$$\|y(t_2)\|_H^2 - \|y(t_1)\|_H^2 + \alpha_1 \int_{t_1}^{t_2} \|y(t)\|_V^p dt \leq \int_{t_1}^{t_2} c_1(t + \tau + h) dt, \tag{12}$$

for each  $t_2 \geq t_1 \geq 0$ , where  $c_1(\cdot)$  is t.u.i. in  $L_1^{\text{loc}}(\mathbb{R}_+)$ . Since the embedding  $V \subset H$  is compact, then this embedding is continuous. So, there exists a constant  $\beta > 0$  such that  $\|b\|_H \leq \beta \|b\|_V$  for each  $b \in V$ . According to (12), since the inequality  $a^2 \leq 1 + a^p$  holds for each  $a \geq 0$ , then the following inequality holds:

$$\|y(t_2)\|_H^2 - \|y(t_1)\|_H^2 + \alpha_3 \int_{t_1}^{t_2} \|y(t)\|_H^2 dt \leq \int_{t_1}^{t_2} [c_1(t + \tau + h) + \alpha_3] dt, \quad (13)$$

for each  $t_2 \geq t_1 \geq 0$ , where  $\alpha_3 = \frac{\alpha_1}{\beta^p}$ . Let us set

$$\rho(t) := \|y(t)\|_H^2 + \alpha_3 \int_0^t \|y(s)\|_H^2 ds - \int_0^t [c_1(s + \tau + h) + \alpha_3] ds, \quad t \geq 0.$$

Inequality (13) and Ball [3, Lemma 7.1] yield that  $\frac{d}{dt}\rho \leq 0$  in  $D^*((0, +\infty))$ , where  $\frac{d}{dt}$  is the derivative operation in the sense of  $D^*((0, +\infty))$ . Thus,

$$\frac{d}{dt} [\|y(t)\|_H^2 + \alpha_3 \|y(t)\|_H^2 - [c_1(t + \tau + h) + \alpha_3] \leq 0 \text{ in } D^*((0, +\infty)).$$

Therefore,

$$\frac{d}{dt} [\|y(t)\|_H^2 e^{\alpha_3 t}] - e^{\alpha_3 t} [c_1(t + \tau + h) + \alpha_3] \leq 0 \text{ in } D^*((0, +\infty)). \quad (14)$$

Ball [3, Lemma 7.1] and inequality (14) imply

$$\|y(t_2)\|_H^2 \leq \|y(t_1)\|_H^2 e^{-\alpha_3(t_2-t_1)} + \int_{t_1}^{t_2} e^{-\alpha_3(t_2-t)} [c_1(t + \tau + h) + \alpha_3] dt, \quad (15)$$

for each  $t_2 \geq t_1 \geq 0$ . Therefore,

$$\begin{aligned} \|y(t_2)\|_H^2 &\leq \|y(t_1)\|_H^2 e^{-\alpha_3(t_2-t_1)} + \int_{t_1}^{t_2} e^{-\alpha_3(t_2-t)} [c_1(t + \tau + h) + \alpha_3] dt \leq \\ &\|y(t_1)\|_H^2 e^{-\alpha_3(t_2-t_1)} + 1 + \int_{t_1+\tau+h}^{t_2+\tau+h} e^{-\alpha_3(t_2-t+\tau+h)} c_1(t) dt \leq \\ &\|y(t_1)\|_H^2 e^{-\alpha_3(t_2-t_1)} + 1 + \frac{K}{\alpha_3} + \\ &\int_{t_1+\tau+h}^{t_2+\tau+h} e^{-\alpha_3(t_2-t+\tau+h)} |c_1(t)| \mathbf{I}\{|c_1(t)| \geq K\} dt, \end{aligned}$$

for each  $K > 0, t_2 \geq t_1 \geq 0$ . Since the function  $c_1 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is t.u.i. in  $L_1^{\text{loc}}(\mathbb{R}_+)$  (see assumption (A1)), then there exists  $K_0 > 0$  such that

$$\sup_{t \geq 0} \int_t^{t+1} |c_1(s)| \mathbf{I}\{|c_1(s)| \geq K_0\} ds \leq 1.$$

Thus,

$$\|y(t_2)\|_H^2 \leq \|y(t_1)\|_H^2 e^{-\alpha_3(t_2-t_1)} + 1 + \frac{K_0}{\alpha_3} + e^{\alpha_3} + 1,$$

that yields estimate (10) with  $c_3 := \alpha_3$  and  $c_4 := 1 + \frac{K_0}{\alpha_3} + e^{\alpha_3} + 1$ , where the positive constants  $c_3$  and  $c_4$  do not depend on  $y \in \mathcal{K}^+$  and  $t \geq 0$ .

Let us prove statement (b). Let  $\{y_n\}_{n \geq 1} \subset \mathcal{K}^+$  be an arbitrary sequence that is bounded in  $L_\infty(\mathbb{R}_+; H)$ . Since  $\mathcal{K}_\cup^+$  is the dense set in a Polish space  $\mathcal{K}^+$  endowed

with the topology induced from  $C^{loc}(\mathbb{R}_+; H)$ , then for each  $n \geq 1$  there exists  $u_n \in \mathcal{K}_\cup^+$  such that

$$\rho_{C^{loc}(\mathbb{R}_+; H)}(y_n, u_n) \leq \frac{1}{n}, \text{ for each } n \geq 1. \tag{16}$$

Note that a priori estimate (10) provides that the sequence  $\{u_n\}_{n \geq 1}$  is bounded in  $L_\infty(\mathbb{R}_+; H)$ . Therefore, the rest of the proof establishes statement (b) for the sequence  $\{u_n\}_{n \geq 1}$ .

Let us fix  $n \geq 1$ . Formula (4) provides the existence of  $\tau_n, h_n \geq 0$  and  $z_n(\cdot) \in \mathcal{K}_{\tau_n}^+$  such that  $u_n(\cdot) = z_n(\cdot + \tau_n + h_n)$ . Then, assumptions (A1) and (A2) yield

$$\begin{aligned} \|u_n(t_2)\|_H^2 - \|u_n(t_1)\|_H^2 + \alpha_1 \int_{t_1}^{t_2} \|u_n(t)\|_V^p dt &\leq \int_{t_1}^{t_2} c_1(t + \tau_n + h_n) dt, \\ \int_{t_1}^{t_2} \|u_n'(t)\|_{V^*}^q dt &\leq \alpha_2 \int_{t_1}^{t_2} \|u_n(t)\|_V^p dt + \int_{t_1}^{t_2} c_2(t + \tau_n + h_n) dt, \end{aligned} \tag{17}$$

for each  $t_2 \geq t_1 \geq 0$  and  $n \geq 1$ .

We remark that

$$\sup_{n \geq 1} \int_{t_1}^{t_2} |c_1(t + \tau_n + h_n)| dt < \infty \text{ and } \sup_{n \geq 1} \int_{t_1}^{t_2} |c_2(t + \tau_n + h_n)| dt < \infty, \tag{18}$$

for each  $t_2 \geq t_1 \geq 0$ , since the functions  $c_1, c_2 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  are t.u.i. in  $L_1^{loc}(\mathbb{R}_+)$ .

Formulae (17) and (18) imply that the sequence  $\{u_n\}_{n \geq 1}$  is bounded in  $W^{loc}(\mathbb{R}_+)$ . Thus, Banach–Alaoglu theorem and Zgurovsky et al. [44, Theorems 1.16 and 1.21] yield that there exist an increasing sequence  $\{n_k\}_{k \geq 1} \subseteq \mathbb{N}$  and elements  $y \in W^{loc}(\mathbb{R}_+) \subset C^{loc}(\mathbb{R}_+; H)$  and  $\bar{c}_1 \in L_1^{loc}(\mathbb{R}_+)$  such that

$$\begin{aligned} u_{n_k} &\rightarrow y && \text{weakly in } L_p^{loc}(\mathbb{R}_+; V), \\ u'_{n_k} &\rightarrow y' && \text{weakly in } L_q^{loc}(\mathbb{R}_+; V^*), \\ u_{n_k} &\rightarrow y && \text{weakly in } C^{loc}(\mathbb{R}_+; H), \\ u_{n_k}(t) &\rightarrow y(t) && \text{in } H \text{ for a.e. } t > 0, \\ c_1(\cdot + \tau_{n_k} + h_{n_k}) &\rightarrow \bar{c}_1 && \text{weakly in } L_1^{loc}(\mathbb{R}_+), \quad k \rightarrow \infty, \end{aligned} \tag{19}$$

where the last convergence holds due to the fact that  $c_1 \in L_1^{loc}(\mathbb{R}_+)$  is t.u.i. in  $L_1^{loc}(\mathbb{R}_+)$ . According to (19), we can pass to the limit in (2). So, we obtain that  $y$  satisfies (2).

We consider the continuous and nonincreasing (by assumption (A1)) functions on  $\mathbb{R}_+$ :

$$\begin{aligned} J_k(t) &= \|u_{n_k}(t)\|_H^2 - \int_0^t c_1(s + \tau_{n_k} + h_{n_k}) ds, \\ J(t) &= \|y(t)\|_H^2 - \int_0^t \bar{c}_1(s) ds, \quad k \geq 1; \end{aligned} \tag{20}$$

cf. Kapustyan and Valero et al. [19]. The last two statements in (19) imply

$$J_k(t) \rightarrow J(t), \text{ as } k \rightarrow +\infty, \text{ for a.e. } t > 0. \tag{21}$$

Similarly to Zgurovsky et al. [46, p. 57] (see the book and references therein) we show that (11) holds. By contradiction suppose the existence of a positive constant  $L > 0$ , a finite interval  $[\tau, T] \subset (0, +\infty)$ , and a subsequence  $\{u_{k_j}\}_{j \geq 1} \subseteq \{u_{n_k}\}_{k \geq 1}$  such that

$$\max_{t \in [\tau, T]} \|u_{k_j}(t) - y(t)\|_H = \|u_{k_j}(t_j) - y(t_j)\|_H \geq L,$$

for each  $j \geq 1$ . Suppose also that  $t_j \rightarrow t_0 \in [\tau, T]$ , as  $j \rightarrow +\infty$ . Continuity of  $\Pi_{\tau, T}y : [\tau, T] \rightarrow H$  implies

$$\liminf_{j \rightarrow +\infty} \|u_{k_j}(t_j) - y(t_0)\|_H \geq L. \tag{22}$$

On the other hand, we prove that

$$u_{k_j}(t_j) \rightarrow y(t_0) \text{ in } H, \quad j \rightarrow +\infty. \tag{23}$$

For this purpose we firstly note that from (19) we have

$$u_{k_j}(t_j) \rightarrow y(t_0) \text{ weakly in } H, \quad j \rightarrow +\infty. \tag{24}$$

Secondly we prove that

$$\limsup_{j \rightarrow +\infty} \|u_{k_j}(t_j)\|_H \leq \|y(t_0)\|_H. \tag{25}$$

We consider the continuous nonincreasing functions  $J$  and  $J_{k_j}$ ,  $j \geq 1$ , defined in (20). Let us fix an arbitrary  $\varepsilon > 0$ . The continuity of  $J$  and (21) provide the existence of  $\bar{t} \in (\tau, t_0)$  such that  $\lim_{j \rightarrow \infty} J_{k_j}(\bar{t}) = J(\bar{t})$  and  $|J(\bar{t}) - J(t_0)| < \varepsilon$ . Then,

$$J_{k_j}(t_j) - J(t_0) \leq |J_{k_j}(\bar{t}) - J(\bar{t})| + |J(\bar{t}) - J(t_0)| \leq |J_{k_j}(\bar{t}) - J(\bar{t})| + \varepsilon,$$

for rather large  $j \geq 1$ . Thus,  $\limsup_{j \rightarrow +\infty} J_{k_j}(t_j) \leq J(t_0)$  and inequality (25) holds.

Thirdly note that the convergence (23) holds due to (24), (25); cf. Gajewski et al. [11, Chapter I]. Finally, we remark that statement (23) contradicts assumption (22). Therefore, according to (16), the first statement of the theorem holds for each sequence  $\{y_n\}_{n \geq 1} \subset \mathcal{K}^+$ .

To finish the proof of statement (b) we note that if, additionally, there exists  $y_0 \in H$  such that  $y_{n_k}(0) \rightarrow y_0$  in  $H$ , then, according to the third convergence in (19),  $y(0) = y_0$ . □

Let us provide the proof of the main result.

*Proof of Theorem 2.1.* Theorem 3.1 implies the following properties for the m-semiflow  $G$ , defined in (8):

- (a) for each  $t \geq 0$  the mapping  $G(t, \cdot) : \mathbb{X} \rightarrow 2^{\mathbb{X}} \setminus \{\emptyset\}$  has a closed graph;
- (b) for each  $t \geq 0$  and  $y_0 \in \mathbb{X}$  the set  $G(t, y_0)$  is compact in  $\mathbb{X}$ ;
- (c) the set  $G(1, \tilde{C})$ , where  $\tilde{C} := \{z \in \mathbb{X} : \|z\|_H^2 < c_4 + 1\}$ , is precompact and attracts each bounded subset  $C \subset \mathbb{X}$ .

Indeed, property (a) follows from Theorem 3.1 (see formulae (5) and (8)); property (b) directly follows from (a) and Theorem 3.1(b); property (c) holds, since  $G(1, \tilde{C})$  is precompact in  $\mathbb{X}$  (Theorem 3.1(b) and formula (8)) and the following inequalities and equality hold:

$$\begin{aligned} \text{dist}_{\mathbb{X}}(G(t, C), G(1, \tilde{C})) &\leq \text{dist}_{\mathbb{X}}(G(1, G(t-1, C)), G(1, \tilde{C})) \leq \\ &\text{dist}_{\mathbb{X}}(G(1, \tilde{C}), G(1, \tilde{C})) = 0, \end{aligned}$$

for sufficiently large  $t$ .

According to properties (a)–(c), Mel’nik and Valero [31, Theorems 1, 2, Remark 2, Proposition 1] yields that the m-semiflow  $G$  has a compact global attractor  $\mathfrak{R}$  in the phase space  $\mathbb{X}$ . □

**4. Applications.** In the following three examples we examine the uniform global attractor for the family of solution sets  $\{\mathcal{K}_\tau^+\}$  generated by particular evolution problems. In all the cases we assume that

$$\forall z \in H \quad \forall \tau \geq 0 \quad \exists y \in \mathcal{K}_\tau^+ \text{ such that } y(\tau) = z.$$

This assumption guarantees the equality  $\mathbb{X} = H$ .

**Example 4.1** (Autonomous evolution problem) Let  $\{\mathcal{K}_\tau^+\}$  be a family of solutions for an autonomous problem on  $[\tau, +\infty)$ ,  $\tau \geq 0$ . Then we have:

$$\forall h \geq 0 \quad T(h)\mathcal{K}_0^+ \subset \mathcal{K}_0^+; \tag{26}$$

$$\forall \tau \geq 0 \quad \forall y \in \mathcal{K}_\tau^+ \quad y(\cdot + \tau) \in \mathcal{K}_0^+. \tag{27}$$

So,  $\mathcal{K}_\cup^+ = \mathcal{K}_0^+$ . If additionally we have that

$$\mathcal{K}_0^+ \text{ is closed in } C^{\text{loc}}(\mathbb{R}_+; H), \tag{28}$$

then

$$\mathcal{K}^+ = \mathcal{K}_0^+.$$

It implies that the m-semiflow  $G$  (defined by (8)) is a classical multivalued semigroup generated by an autonomous evolution problem.

**Example 4.2** (Non-autonomous evolution problem) Let  $\{\mathcal{K}_\tau^+\}$  be a family of solutions for non-autonomous problem on  $[\tau, +\infty)$ ,  $\tau \geq 0$ , and the following condition holds:

$$\forall s \geq \tau \geq 0 \quad \forall y \in \mathcal{K}_\tau^+ \quad \Pi_{s,+\infty}y(\cdot) \in \mathcal{K}_s^+. \tag{29}$$

Then, according to Kapustyan et al. [21], formula

$$U(t, \tau, z) = \{y(t) : y(\cdot) \in \mathcal{K}_\tau^+, y(\tau) = z\} \tag{30}$$

defines a m-semiprocess, that is

$$\forall t \geq s \geq \tau \quad U(t, \tau, z) \subset U(t, s, U(s, \tau, z)).$$

One of the most important objects for m-semiprocess (30) is uniform global attractor; Chepyzhov and Vishik[7], Kapustyan et al. [20], Zgurovsky et al. [46]. It is a set  $\Theta$  such that for every bounded subset  $C \subset H$

$$\sup_{\tau \geq 0} \text{dist}_H(U(t + \tau, \tau, C), \Theta) \rightarrow 0, \quad t \rightarrow \infty, \tag{31}$$

and  $\Theta$  is minimal among all closed sets satisfying this property. Then under assumptions (A1), (A2) and from (29) it follows that the m-semiprocess (30) has the compact uniform global attractor  $\Theta \subseteq \mathfrak{R}$ , where  $\mathfrak{R}$  is the global attractor for the m-semiflow (8).

Indeed,

$$\forall t \geq \tau \geq 0 \quad \forall z \in H \quad U(t + \tau, \tau, z) \subset G(t, z). \tag{32}$$

So, if  $\mathfrak{R}$  is a compact global attractor for the m-semiflow  $G$  then, according to Kapustyan et al. [20], there exists a compact uniform global attractor  $\Theta$  for m-semiprocess  $U$  and, moreover,  $\Theta \subset \mathfrak{R}$ .

In the following example we examine the existence of uniform global attractor for non-autonomous differential-operator inclusion. The uniform trajectory attractors



for classes of non-autonomous inclusions and equations were proved to exist in Zgurovsky and Kasyanov [48] (see also Gorban et al. [13]).

**Example 4.3** (Non-autonomous differential-operator inclusion) For the multivalued map  $A : \mathbb{R}_+ \times V \rightarrow 2^{V^* \setminus \{\emptyset\}}$  we consider the problem of long-time behavior of all globally defined weak solutions for non-autonomous evolution inclusion

$$y'(t) + A(t, y(t)) \ni \bar{0}, \tag{33}$$

as  $t \rightarrow +\infty$ . Let  $\langle \cdot, \cdot \rangle_V : V^* \times V \rightarrow \mathbb{R}$  be the pairing in  $V^* \times V$ , that coincides on  $H \times V$  with the inner product  $(\cdot, \cdot)$  in the Hilbert space  $H$ .

We note that Problem (33) arises in many important models for distributed parameter control problems and that large class of identification problems enter this formulation. Let us indicate a problem which is one of the motivations for the study of the non-autonomous evolution inclusion (33) (see, for example, Migórski and Ochal [33]; Zgurovsky et al. [46] and references therein). In a subset  $\Omega$  of  $\mathbb{R}^3$ , we consider the nonstationary heat conduction equation

$$\frac{\partial y}{\partial t} - \Delta y = f \text{ in } \Omega \times (0, +\infty)$$

with initial conditions and suitable boundary ones. Here  $y = y(x, t)$  represents the temperature at the point  $x \in \Omega$  and time  $t > 0$ . It is supposed that  $f = f_1 + f_2$ , where  $f_2$  is given and  $f_1$  is a known function of the temperature of the form

$$-f_1(x, t) \in \partial j(x, t, y(x, t)) \text{ a.e. } (x, t) \in \Omega \times (0, +\infty).$$

Here  $\partial j(x, t, \xi)$  denotes generalized gradient of Clarke (see Clarke [9]) with respect to the last variable of a function  $j : \Omega \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$  which is assumed to be locally Lipschitz in  $\xi$  (cf. Migórski and Ochal [33] and references therein). The multivalued function  $\partial j(x, t, \cdot) : \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is generally nonmonotone and it includes the vertical jumps. In a physicist’s language it means that the law is characterized by the generalized gradient of a nonsmooth potential  $j$  (cf. Panagiotopoulos [35]). Models of physical interest includes also the next (see, for example, Balibrea et al. [2] and references therein): a model of combustion in porous media; a model of conduction of electrical impulses in nerve axons; a climate energy balance model; etc.

Let the following assumptions hold:

- (H1) (*Growth condition*) There exist a t.u.i. in  $L_1^{\text{loc}}(\mathbb{R}_+)$  function  $c_1 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  and a constant  $c_2 > 0$  such that  $\|d\|_{V^*}^q \leq c_1(t) + c_2\|u\|_V^p$  for any  $u \in V$ ,  $d \in A(t, u)$ , and a.e.  $t > 0$ ;
- (H2) (*Sign condition*) There exist a constant  $\alpha > 0$  and a t.u.i. in  $L_1^{\text{loc}}(\mathbb{R}_+)$  function  $\beta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that  $\langle d, u \rangle_V \geq \alpha\|u\|_V^p - \beta(t)$  for any  $u \in V$ ,  $d \in A(t, u)$ , and a.e.  $t > 0$ ;
- (H3) (*Strong measurability*) If  $C \subseteq V^*$  is a closed set, then the set  $\{(t, u) \in (0, +\infty) \times V : A(t, u) \cap C \neq \emptyset\}$  is a Borel subset in  $(0, +\infty) \times V$ ;
- (H4) (*Pointwise pseudomonotonicity*) Let for a.e.  $t > 0$  the following two assumptions hold:
  - a) for every  $u \in V$  the set  $A(t, u)$  is nonempty, convex, and weakly compact one in  $V^*$ ;
  - b) if a sequence  $\{u_n\}_{n \geq 1}$  converges weakly in  $V$  towards  $u \in V$  as  $n \rightarrow +\infty$ ,  $d_n \in A(t, u_n)$  for any  $n \geq 1$ , and  $\limsup_{n \rightarrow +\infty} \langle d_n, u_n - u \rangle_V \leq 0$ ,

then for any  $\omega \in V$  there exists  $d(\omega) \in A(t, u)$  such that

$$\liminf_{n \rightarrow +\infty} \langle d_n, u_n - \omega \rangle_V \geq \langle d(\omega), u - \omega \rangle_V.$$

Let  $0 \leq \tau < T < +\infty$ . As a *weak solution* of evolution inclusion (33) on the interval  $[\tau, T]$  we consider an element  $u(\cdot)$  of the space  $L_p(\tau, T; V)$  such that for some  $d(\cdot) \in L_q(\tau, T; V^*)$  it is fulfilled:

$$-\int_{\tau}^T (\xi'(t), y(t)) dt + \int_{\tau}^T \langle d(t), \xi(t) \rangle_V dt = 0 \quad \forall \xi \in C_0^\infty([\tau, T]; V), \quad (34)$$

and  $d(t) \in A(t, y(t))$  for a.e.  $t \in (\tau, T)$ . For fixed nonnegative  $\tau$  and  $T$ ,  $\tau < T$ , let us consider

$$X_{\tau, T} = L_p(\tau, T; V), \quad X_{\tau, T}^* = L_q(\tau, T; V^*),$$

$$W_{\tau, T} = \{y \in X_{\tau, T} \mid y' \in X_{\tau, T}^*\}, \quad \mathcal{A}_{\tau, T} : X_{\tau, T} \rightarrow 2^{X_{\tau, T}^*} \setminus \{\emptyset\},$$

$$\mathcal{A}_{\tau, T}(y) = \{d \in X_{\tau, T}^* \mid d(t) \in A(t, y(t)) \text{ for a.e. } t \in (\tau, T)\},$$

where  $y'$  is a derivative of an element  $u \in X_{\tau, T}$  in the sense of  $\mathcal{D}([\tau, T]; V^*)$  (see, for example, Gajewski, Gröger, and Zacharias [11, Definition IV.1.10]). Gajewski, Gröger, and Zacharias [11, Theorem IV.1.17] provide that the embedding  $W_{\tau, T} \subset C([\tau, T]; H)$  is continuous and dense. Moreover,

$$(u(T), v(T)) - (u(\tau), v(\tau)) = \int_{\tau}^T [\langle u'(t), v(t) \rangle_V + \langle v'(t), u(t) \rangle_V] dt, \quad (35)$$

for any  $u, v \in W_{\tau, T}$ .

Migórski [34, Lemma 7, p. 516] (see the paper and references therein) and the assumptions above provide that the multivalued mapping  $\mathcal{A}_{\tau, T} : X_{\tau, T} \rightarrow 2^{X_{\tau, T}^*} \setminus \{\emptyset\}$  satisfies the listed below properties:

- (P1) There exists a positive constant  $C_1 = C_1(\tau, T)$  such that  $\|d\|_{X_{\tau, T}^*} \leq C_1(1 + \|y\|_{X_{\tau, T}}^{p-1})$  for any  $y \in X_{\tau, T}$  and  $d \in \mathcal{A}_{\tau, T}(y)$ ;
- (P2) There exist positive constants  $C_2 = C_2(\tau, T)$  and  $C_3 = C_3(\tau, T)$  such that  $\langle d, y \rangle_{X_{\tau, T}} \geq C_2 \|y\|_{X_{\tau, T}}^p - C_3$  for any  $y \in X_{\tau, T}$  and  $d \in \mathcal{A}_{\tau, T}(y)$ ;
- (P3) The multivalued mapping  $\mathcal{A}_{\tau, T} : X_{\tau, T} \rightarrow 2^{X_{\tau, T}^*} \setminus \{\emptyset\}$  is (generalized) pseudomonotone on  $W_{\tau, T}$ , i.e.
  - a) for every  $y \in X_{\tau, T}$  the set  $\mathcal{A}_{\tau, T}(y)$  is a nonempty, convex and weakly compact one in  $X_{\tau, T}^*$ ;
  - b)  $\mathcal{A}_{\tau, T}$  is upper semi-continuous from every finite dimensional subspace  $X_{\tau, T}$  into  $X_{\tau, T}^*$  endowed with the weak topology;
  - c) if a sequence  $\{y_n, d_n\}_{n \geq 1} \subset W_{\tau, T} \times X_{\tau, T}^*$  converges weakly in  $W_{\tau, T} \times X_{\tau, T}^*$  towards  $(y, d) \in W_{\tau, T} \times X_{\tau, T}^*$ ,  $d_n \in \mathcal{A}_{\tau, T}(y_n)$  for any  $n \geq 1$ , and  $\limsup_{n \rightarrow +\infty} \langle d_n, y_n - y \rangle_{X_{\tau, T}} \leq 0$ , then  $d \in \mathcal{A}_{\tau, T}(y)$  and  $\lim_{n \rightarrow +\infty} \langle d_n, y_n \rangle_{X_{\tau, T}} = \langle d, y \rangle_{X_{\tau, T}}$ .

Formula (34) and the definition of the derivative for an element from  $\mathcal{D}([\tau, T]; V^*)$  yield that each weak solution  $y \in X_{\tau, T}$  of Problem (33) on  $[\tau, T]$  belongs to the space  $W_{\tau, T}$  and  $y' + \mathcal{A}_{\tau, T}(y) \ni \bar{0}$ . On the contrary, suppose that  $y \in W_{\tau, T}$  satisfies the last inclusion, then  $y$  is a weak solution of Problem (33) on  $[\tau, T]$ .

Assumption (H1), properties (P1)–(P3), and Denkowski, Migórski, and Papa-georgiou [10, Theorem 1.3.73] (see also Zgurovsky, Mel'nik, and Kasyanov [44, Chapter 2] and references therein) provide the existence of a weak solution of Cauchy problem (33) with initial data  $y(\tau) = y^{(\tau)}$  on the interval  $[\tau, T]$ , for any  $y^{(\tau)} \in H$ .

For fixed  $\tau$  and  $T$ , such that  $0 \leq \tau < T < +\infty$ , we denote

$$\mathcal{D}_{\tau,T}(y^{(\tau)}) = \{y(\cdot) \mid y \text{ is a weak solution of (33) on } [\tau, T], y(\tau) = y^{(\tau)}\}, \quad y^{(\tau)} \in H.$$

We remark that  $\mathcal{D}_{\tau,T}(y^{(\tau)}) \neq \emptyset$  and  $\mathcal{D}_{\tau,T}(y^{(\tau)}) \subset W_{\tau,T}$ , if  $0 \leq \tau < T < +\infty$  and  $y^{(\tau)} \in H$ . Moreover, the concatenation of weak solutions of Problem (33) is a weak solutions too, i.e. if  $0 \leq \tau < t < T$ ,  $y^{(\tau)} \in H$ ,  $y(\cdot) \in \mathcal{D}_{\tau,t}(y^{(\tau)})$ , and  $v(\cdot) \in \mathcal{D}_{t,T}(y(t))$ , then

$$z(s) = \begin{cases} y(s), & s \in [\tau, t], \\ v(s), & s \in [t, T], \end{cases}$$

belongs to  $\mathcal{D}_{\tau,T}(y^{(\tau)})$ ; cf. Zgurovsky et al. [46, pp. 55–56].

Gronwall's lemma provides that for any finite time interval  $[\tau, T] \subset \mathbb{R}_+$  each weak solution  $y$  of Problem (33) on  $[\tau, T]$  satisfies the estimates

$$\|y(t)\|_H^2 - 2 \int_0^t \beta(\xi) d\xi + 2\alpha \int_s^t \|y(\xi)\|_V^p d\xi \leq \|y(s)\|_H^2 - 2 \int_0^s \beta(\xi) d\xi, \quad (36)$$

$$\|y(t)\|_H^2 \leq \|y(s)\|_H^2 e^{-2\alpha\gamma(t-s)} + 2 \int_s^t (\beta(\xi) + \alpha\gamma) e^{-2\alpha\gamma(t-\xi)} d\xi, \quad (37)$$

where  $t, s \in [\tau, T]$ ,  $t \geq s$ ;  $\gamma > 0$  is a constant such that  $\gamma \|u\|_H^p \leq \|u\|_V^p$  for any  $u \in V$ ; cf. Zgurovsky et al. [46, p. 56]. In the proof of (37) we used the inequality  $\|u\|_H^2 - 1 \leq \|u\|_H^p$  for any  $u \in H$ .

Therefore, any weak solution  $y$  of Problem (33) on a finite time interval  $[\tau, T] \subset \mathbb{R}_+$  can be extended to a global one, defined on  $[\tau, +\infty)$ . For arbitrary  $\tau \geq 0$  and  $y^{(\tau)} \in H$  let  $\mathcal{D}_\tau(y^{(\tau)})$  be the set of all weak solutions (defined on  $[\tau, +\infty)$ ) of Problem (33) with initial data  $y(\tau) = y^{(\tau)}$ . Let us consider the family  $\mathcal{K}_\tau^+ = \cup_{y^{(\tau)} \in H} \mathcal{D}_\tau(y^{(\tau)})$  of all weak solutions of Problem (33) defined on the semi-infinite time interval  $[\tau, +\infty)$ .

Properties (P1)–(P2) imply assumptions (A1) and (A2). Therefore, Theorem 2.1 yields that the m-semiflow  $G$ , defined in (8), has a compact global attractor  $\mathfrak{R}$  in the phase space  $H$ .

**Acknowledgments.** Authors thank Prof. Oleksiy V. Kapustyan for useful discussions during the preparation of this paper. Authors thank the anonymous referees for their constructive suggestions and remarks. The research was partially supported by the National Academy of Sciences of Ukraine under grant 2284/15 and by Grant of the President of Ukraine GP/F61/017.

REFERENCES

[1] A. V. Babin and M. I. Vishik, *Attractors of Evolution Equations* (Russian), Nauka, Moscow, 1989.

[2] F. Balibrea, T. Caraballo, P. E. Kloeden and J. Valero, [Recent developments in dynamical systems: Three perspectives](#), *International Journal of Bifurcation and Chaos*, 20 (2010), 2591–2636.

[3] J. M. Ball, [Continuity properties and global attractors of generalized semiflows and the Navier-Stokes equations](#). *Nonlinear Science*, 7 (1997), 475–502 Erratum, *ibid* 8:233,1998. Corrected version appears in *Mechanics: From Theory to Computation*. Springer Verlag, (2000), 447–474.

[4] V. V. Chepyzhov and M. I. Vishik, [Trajectory attractors for evolution equations](#), *C.R.Acad.Sci. Paris. Serie I*, 321 (1995), 1309–1314.

[5] V. V. Chepyzhov and M. I. Vishik, [Evolution equations and their trajectory attractors](#), *J. Math. Pures Appl.*, 76 (1997), 913–964.

- [6] V. V. Chepyzhov and M. I. Vishik, [Trajectory and global attractors for 3D Navier-Stokes system](#), *Mat. Zametki*, **71** (2002), 177–193.
- [7] V. V. Chepyzhov and M. I. Vishik, *Attractors for Equations of Mathematical Physics*, American Mathematical Society, Providence RI, 2002.
- [8] V. V. Chepyzhov and M. I. Vishik, [Trajectory attractor for reaction-diffusion system with diffusion coefficient vanishing in time](#), *Discrete and Continuous Dynamical Systems*, **27** (2010), 1498–1509.
- [9] F. H. Clarke, *Optimization and Nonsmooth Analysis*. John Wiley & Sons, Inc., New York, 1983.
- [10] Z. Denkowski, S. Migórski and N. S. Papageorgiou, *An Introduction to Nonlinear Analysis: Applications*, Kluwer Academic/Plenum Publishers, Boston, 2003.
- [11] H. Gajewski, K. Gröger and K. Zacharias, *Nichtlineare Operatorgleichungen und Operatordifferentialgleichungen*, Akademie-Verlag, Berlin, 1975.
- [12] M. O. Gluzman, N. V. Gorban and P. O. Kasyanov, [Lyapunov type functions for classes of autonomous parabolic feedback control problems and applications](#), *Applied Mathematics Letters*, **39** (2015), 19–21.
- [13] N. V. Gorban, O. V. Kapustyan and P. O. Kasyanov, [Uniform trajectory attractor for non-autonomous reaction-diffusion equations with Caratheodory’s nonlinearity](#), *Nonlinear Analysis, Theory, Methods and Applications*, **98** (2014), 13–26.
- [14] N. V. Gorban, O. V. Kapustyan, P. O. Kasyanov and L. S. Paliichuk, [On global attractors for autonomous damped wave equation with discontinuous nonlinearity](#), *Continuous and Distributed Systems. Theory and Applications, Solid Mechanics and its Applications*, **211** (2014), 221–237.
- [15] N. V. Gorban and P. O. Kasyanov, [On regularity of all weak solutions and their attractors for reaction-diffusion inclusion in unbounded domain](#), *Continuous and Distributed Systems. Theory and Applications, Solid Mechanics and its Applications*, **211** (2013), 205–220.
- [16] J. K. Hale, *Asymptotic Behavior of Dissipative Systems*, AMS, Providence, RI, 1988.
- [17] G. Iovane, A. V. Kapustyan and J. Valero, [Asymptotic behavior of reaction-diffusion equations with non-damped impulsive effects](#), *Nonlinear Analysis*, **68** (2008), 2516–2530.
- [18] A. V. Kapustyan and J. Valero, [Weak and strong attractors for the 3D Navier-Stokes system](#), *Journal of Differential Equations*, **240** (2007), 249–278.
- [19] A. V. Kapustyan and J. Valero, [On the Kneser property for the complex Ginzburg-Landau equation and the Lotka-Volterra system with diffusion](#), *J Math Anal Appl.*, **357** (2009), 254–272.
- [20] O. V. Kapustyan, P. O. Kasyanov and J. Valero, [Pullback attractors for a class of extremal solutions of the 3D Navier-Stokes equations](#), *Journal of Mathematical Analysis and Applications*, **373** (2011), 535–547.
- [21] A. V. Kapustyan, P. O. Kasyanov, J. Valero and M. Z. Zgurovsky, [Structure of uniform global attractor for general non-autonomous reaction-diffusion system](#), *Continuous and Distributed Systems: Theory and Applications, Solid Mechanics and Its Applications*, **211** (2014), 163–180.
- [22] O. V. Kapustyan, P. O. Kasyanov and J. Valero, [Structure and regularity of the global attractor of a reaction-diffusion equation with non-smooth nonlinear term](#), *Discrete and Continuous Dynamical Systems, Series A*, **34** (2014), 4155–4182.
- [23] O. V. Kapustyan, P. O. Kasyanov and J. Valero, [Regular solutions and global attractors for reaction-diffusion systems without uniqueness](#), *Communications on Pure and Applied Analysis*, **13** (2014), 1891–1906.
- [24] P. O. Kasyanov, V. S. Mel’nik and S. Toscano, [Solutions of Cauchy and periodic problems for evolution inclusions with multi-valued  \$w\_{\lambda\_0}\$ -pseudomonotone maps](#), *Journal of Differential Equations*, **249** (2010), 1258–1287.
- [25] P. O. Kasyanov, [Multivalued dynamics of solutions of an autonomous differential-operator inclusion with pseudomonotone nonlinearity](#), *Cybernetics and Systems Analysis*, **47** (2011), 800–811.
- [26] P. O. Kasyanov, [Multivalued dynamics of solutions of autonomous operator differential equations with pseudomonotone nonlinearity](#), *Mathematical Notes*, **92** (2012), 205–218.
- [27] P. O. Kasyanov, L. Toscano and N. V. Zadoianchuk, [Regularity of weak solutions and their attractors for a parabolic feedback control problem](#), *Set-Valued and Variational Analysis*, **21** (2013), 271–282.

- [28] P. E. Kloeden, P. Marin-Rubio and J. Valero, [The envelope attractor of non-strict multivalued dynamical systems with application to the 3d navier-stokes and reaction-diffusion equations](#), *Set-Valued and Variational Analysis*, **21** (2013), 517–540.
- [29] M. A. Krasnosel'skii, *Topological Methods in the Theory of Nonlinear Integral Equations*, (International Series of Monographs on Pure and Applied Mathematics, Vol. 45) Oxford/London/New York/Paris, 1964.
- [30] O. A. Ladyzhenskaya, *Attractors for Semigroups and Evolution Equations*, Cambridge University Press, Cambridge, 1991.
- [31] V. S. Melnik and J. Valero, [On attractors of multivalued semi-flows and generalized differential equations](#), *Set-Valued Anal.*, **6** (1998), 83–111.
- [32] V. S. Mel'nik and J. Valero, [On global attractors of multivalued semiprocesses and non-autonomous evolution inclusions](#), *Set-Valued Anal.*, **8** (2000), 375–403.
- [33] S. Migórski and A. Ochal, [Optimal Control of Parabolic Hemivariational Inequalities](#), *Journal of Global Optimization*, **17** (2000), 285–300.
- [34] S. Migórski, [Boundary hemivariational inequalities of hyperbolic type and applications](#), *Journal of Global Optimization*, **31** (2005), 505–533.
- [35] P. D. Panagiotopoulos, *Inequality Problems in Mechanics and Applications*, Convex and Nonconvex Energy Functions, Birkhauser, Basel, 1985.
- [36] G. R. Sell, [Global attractors for the three-dimensional Navier-Stokes equations](#), *J. Dyn. Diff. Eq.*, **8** (1996), 1–33.
- [37] J. Smoller, *Shock Waves and Reaction-Diffusion Equations (Grundlehren der Mathematischen Wissenschaften)*, Springer-Verlag, New York, 1983.
- [38] H. Sohr, *The Navier-Stokes Equations. An Elementary Functional Analytic Approach*, Verlag, Birkhäuser, 2001.
- [39] R. Temam, *Navier-Stokes Equations*, North-Holland, Amsterdam, 1979.
- [40] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, Appl. Math. Sci., Springer-Verlag, New York, 1988.
- [41] J. Valero and A. V. Kapustyan, [On the connectedness and asymptotic behaviour of solutions of reaction-diffusion systems](#), *J Math Anal Appl.*, **323** (2006), 614–633.
- [42] M. I. Vishik, S. V. Zelik and V. V. Chepyzhov, [Strong trajectory attractor for a dissipative reaction-diffusion system](#), *Doklady Mathematics*, **82** (2010), 869–873.
- [43] J. Warga, *Optimal Control of Differential and Functional Equations*, Academic Press, 1972.
- [44] M. Z. Zgurovsky, V. S. Mel'nik and P. O. Kasyanov, *Evolution Inclusions and Variation Inequalities for Earth Data Processing II*, Springer, Berlin, 2011.
- [45] M. Z. Zgurovsky, P. O. Kasyanov and N. V. Zadoianchuk (Zadoyanchuk), [Long-time behavior of solutions for quasilinear hyperbolic hemivariational inequalities with application to piezoelectricity problem](#), *Applied Mathematics Letters*, **25** (2012), 1569–1574.
- [46] M. Z. Zgurovsky, P. O. Kasyanov, O. V. Kapustyan, J. Valero and N. V. Zadoianchuk, *Evolution Inclusions and Variation Inequalities for Earth Data Processing III*, Springer, Berlin, 2012.
- [47] M. Z. Zgurovsky and P. O. Kasyanov, [Multivalued dynamics of solutions for autonomous operator differential equations in strongest topologies](#), *Continuous and Distributed Systems. Theory and Applications. Solid Mechanics and its Applications*, **211** (2014), 149–162.
- [48] M. Z. Zgurovsky and P. O. Kasyanov, [Evolution inclusions in Nonsmooth systems with applications for Earth data processing: Uniform trajectory attractors for non-autonomous evolution inclusions solutions with pointwise pseudomonotone mappings](#), *Advances in Global Optimization, Springer Proceedings in Mathematics and Statistics*, **95** (2015), 283–294.

Received January 2016; revised February 2016.

E-mail address: [zgurovsm@hotmail.com](mailto:zgurovsm@hotmail.com)

E-mail address: [mark.gluzman@columbia.edu](mailto:mark.gluzman@columbia.edu)

E-mail address: [nata\\_gorban@i.ua](mailto:nata_gorban@i.ua)

E-mail address: [kasyanov@i.ua](mailto:kasyanov@i.ua)

E-mail address: [lili262808@gmail.com](mailto:lili262808@gmail.com)

E-mail address: [olgkhomenko@ukr.net](mailto:olgkhomenko@ukr.net)